

**BOTSCHAFT
DER REPUBLIK TÜRKEI**

Rungestr. 9, 10179 Berlin
Tel: +49(30)275850 Fax: +49(30)27590915

Az.: 454.60/2007/Berlin BE/34310

Berlin, den 06.12.2007

An das
Bundesministerium für Ernährung, Landwirtschaft
und Verbraucherschutz
Frau Christiane Neikes
Referat 517
- Pflanzenschutz-
Rochusstr. 1

53123 Bonn

071062733

| |
|---|
| Bundesministerium für Ernährung, Landwirtschaft und Verbraucherschutz - Posteingang - |
| Eing. 10. Dez. 2007 |
| Abt./ Ref 5/15/1517 |
| Az.: 3332/42 |

M/12

Sehr geehrte Frau Neikes,

das Ministerium für Landwirtschaft der Republik Türkei teilte mit, dass die Gesundheitskriterien, die gemäß Artikel 5 der Landwirtschaftlichen Quarantäneverordnung die Einfuhr zur Saat geeigneter Kartoffeln betreffen, umgestaltet wurden. Anbei befindet sich diesbezüglich eine detaillierte Beschreibung der umgestalteten Kriterien.

Mit freundlichen Grüßen

Vakur Eysel
Gesandter



Anlage

Fr. Neikes
H. d. V.
M/12

Qualifications required in seed potatoes for some potato diseases which are known as quality diseases in the framework of Plant Quarantine Regulation of Article 5 are as follows:

2% of seed potato consignments are inspected. Sample as 5% of inspected material is taken and examine in respect of quality diseases by inspectors taking into consideration the following standards.

Furthermore 200 tubers for per 100 tones are sent to the laboratory on to be examined in terms of quarantine diseases.

1. Powdery Scab , (*Spongospora subterranea*)

The weight of potato tubers of which a third and more of tuber surface is covered by disease symptoms can not exceed 5% of sample weight.

2. Common Scab (*Streptomyces spp*)

The weight of potato tubers of which a third and more of tuber surface is covered by disease symptoms can not exceed 5% of sample weight.

3. Black Scurf (*Rhizoctonia solani*)

The weight of potato tubers of which a 10% and more of tuber surface is covered by disease symptoms can not exceed 8% of sample weight.

4. Silver Scurf (*Helminthosporium solani*)

The weight of tubers which are started to shrink(=loose turgor) by reason of silver scurf and affect at least one eye can not exceed 5% of sample weight

5. Dry Rot (*Fusarium spp.*):

The weight of tubers which are contaminated with dry rot disease (*Fusarium spp.*) can not exceed 2% of sample weight.

6. Late Blight (*Phytophthora infestans*):

The weight of tubers which show the symptoms of late blight (*Phytophthora infestans*) can not exceed 6% of sample weight.